

5 A third determinant of hybridization efficiency is the salt concentration of the reaction. In large part, the choice of solution conditions will depend on the requirements of the cleavage agent, and for reagents obtained commercially, the manufacturer's instructions are a resource for this information. When developing an assay utilizing any particular cleavage agent, the oligonucleotide and temperature optimizations described above should be performed in the buffer conditions best suited to that cleavage agent.

10 A "no enzyme" control allows the assessment of the stability of the labeled oligonucleotides under particular reaction conditions, or in the presence of the sample to be tested (*i.e.*, in assessing the sample for contaminating nucleases). In this manner, the substrate and oligonucleotides are placed in a tube containing all reaction components, except the enzyme and treated the same as the enzyme-containing reactions. Other controls may also be included. For example, a reaction with all of the components except the target nucleic acid will serve to confirm the dependence of the cleavage on the presence of the target sequence.

15 **Probing For Multiple Alleles**

20 The invader-directed cleavage reaction is also useful in the detection and quantification of individual variants or alleles in a mixed sample population. By way of example, such a need exists in the analysis of tumor material for mutations in genes associated with cancers. Biopsy material from a tumor can have a significant complement of normal cells, so it is desirable to detect mutations even when present in fewer than 5% of the copies of the target nucleic acid in a sample. In this case, it is also desirable to measure what fraction of the population carries the mutation. Similar analyses may also be done to examine allelic variation in other gene systems, and it is not intended that the method of the present invention be limited to the analysis of tumors.

25 As demonstrated below, reactions can be performed under conditions that prevent the cleavage of probes bearing even a single-nucleotide difference mismatch within the region of the target nucleic acid termed "Z" in Figure 29, but that permit

cleavage of a similar probe that is completely complementary to the target in this region. Thus, the assay may be used to quantitate individual variants or alleles within a mixed sample.

The use of multiple, differently labelled probes in such an assay is also contemplated. To assess the representation of different variants or alleles in a sample, one would provide a mixture of probes such that each allele or variant to be detected would have a specific probe (*i.e.*, perfectly matched to the Z region of the target sequence) with a unique label (*e.g.*, no two variant probes with the same label would be used in a single reaction). These probes would be characterized in advance to ensure that under a single set of reaction conditions, they could be made to give the same rate of signal accumulation when mixed with their respective target nucleic acids. Assembly of a cleavage reaction comprising the mixed probe set, a corresponding invader oligonucleotide, the target nucleic acid sample, and the appropriate cleavage agent, along with performance of the cleavage reaction under conditions such that only the matched probes would cleave, would allow independent quantification of each of the species present, and would therefore indicate their relative representation in the target sample.

IV. A Comparision Of Invasive Cleavage And Primer-Directed Cleavage

As discussed herein, the terms "invasive" or "invader-directed" cleavage specifically denote the use of a first, upstream oligonucleotide, as defined below, to cause specific cleavage at a site within a second, downstream sequence. To effect such a direction of cleavage to a region within a duplex, it is required that the first and second oligonucleotides overlap in sequence. That is to say, a portion of the upstream oligonucleotide, termed the "invader", has significant homology to a portion of the downstream "probe" oligonucleotide, so that these regions would tend to basepair with the same complementary region of the target nucleic acid to be detected. While not limiting the present invention to any particular mechanism, the overlapping regions would be expected to alternate in their occupation of the shared hybridization site. When the probe oligonucleotide fully anneals to the target nucleic acid, and thus forces

the 3' region of the invader to remain unpaired, the structure so formed is not a substrate for the 5' nucleases of the present invention. By contrast, when the inverse is true, the structure so formed is substrate for these enzymes, allowing cleavage and release of the portion of the probe oligonucleotide that is displaced by the invader oligonucleotide. The shifting of the cleavage site to a region the probe oligonucleotide that would otherwise be basepaired to the target sequence is one hallmark of the invasive cleavage assay (*i.e.*, the invader-directed cleavage assay) of the present invention.

It is beneficial at this point to contrast the invasive cleavage as described above with two other forms of probe cleavage that may lead to internal cleavage of a probe oligonucleotide, but which do not comprise invasive cleavage. In the first case, a hybridized probe may be subject to duplex-dependent 5' to 3' exonuclease "nibbling," such that the oligonucleotide is shortened from the 5' end until it cannot remain bound to the target (*see, e.g.*, Examples 6-8 and Figs. 26-28). The site at which such nibbling stops can appear to be discrete, and, depending on the difference between the melting temperature of the full-length probe and the temperature of the reaction, this stopping point may be 1 or several nucleotides into the probe oligonucleotide sequence. Such "nibbling" is often indicated by the presence of a "ladder" of longer products ascending size up to that of the full length of the probe, but this is not always the case. While any one of the products of such a nibbling reaction may be made to match in size and cleavage site the products of an invasive cleavage reaction, the creation of these nibbling products would be highly dependent on the temperature of the reaction and the nature of the cleavage agent, but would be independent of the action of an upstream oligonucleotide, and thus could not be construed to involve invasive cleavage.

A second cleavage structure that may be considered is one in which a probe oligonucleotide has several regions of complementarity with the target nucleic acid, interspersed with one or more regions or nucleotides of noncomplementarity. These noncomplementary regions may be thought of as "bubbles" within the nucleic acid duplex. As temperature is elevated, the regions of complementarity can be expected to